arXiv:1401.2236v1 [math.CA] 10 Jan 2014

# ON THE COEFFICIENTS OF AN EXPANSION OF $(1+1/x)^x$ RELATED TO CARLEMAN'S INEQUALITY

YUE HU AND CRISTINEL MORTICI

Abstract. In this note, we present new properties for the sequence $b_1 = \frac{1}{2}$, $b_n = \frac{1}{n}(\frac{1}{n+1} - \sum_{k=0}^{n-2} \frac{b_{n-k-1}}{k+2}), n \geq 2$, arising in some refinements of Carleman's inequality. Our results extend some results of Yang [Approximations for constant $e$ and their applications J. Math. Anal. Appl. 262 (2001) 651-659] and Alzer and Berg [some classes of completely monotonic functions Ann. Acad. Sci. Fennicae 27(2002) 445-460].

**AMS Subject Classification:** 26D15, 33B15
**Keywords**: constant $e$; sequence; inequalities; integral representation

## 1. Introduction

The following Carleman inequality [3]

$$\sum_{n=1}^{\infty} (a_1 a_2 \cdots a_n)^{1/n} < e \sum_{n=1}^{\infty} a_n,$$

whenever $a_n \geq 0$, $n = 1, 2, 3, \ldots$, with $0 < \sum_{n=1}^{\infty} a_n < \infty$, has attracted the attention of many authors in the recent past. We refer for example to [4], or to the work of Yang [6], who proved

$$\sum_{n=1}^{\infty} (a_1 a_2 \cdots a_n)^{1/n} < e \sum_{n=1}^{\infty} \left(1 - \sum_{k=1}^{6} \frac{b_k}{(n+1)^k}\right) a_n$$

with $b_1 = 1/2, b_2 = 1/24, b_3 = 1/48, b_4 = 73/5760, b_5 = 11/1280, b_6 = 1945/580608$. In the final part of his paper, Yang [6] conjectured that if

$$\left(1 + \frac{1}{x}\right)^x = e\left(1 - \sum_{n=1}^{\infty} \frac{b_n}{(x+1)^n}\right), \quad x > 0,$$

then $b_n > 0$, $n = 1, 2, 3, \ldots$ .

Later , this conjecture was proved and discussed by Yang [7], Gylletberg and Yan [2], and Yue [8], using the recurrence

$$b_1 = \frac{1}{2}, b_n = \frac{1}{n}(\frac{1}{n+1} - \sum_{k=0}^{n-2} \frac{b_{n-k-1}}{k+2}) \qquad (n = 2, 3, \cdots). \tag{1.1}$$

Here we observe that it is easy to calculate the value of $b_n$ by computer programs, but it is hard to extract any properties about $b_n$ from (1.1). Also the proofs of $b_n > 0$ provided were quite complicated.

1991 *Mathematics Subject Classification.* 26D15; 33B15.
*Key words and phrases.* Carleman inequality; integral representation; series.

In this note, we use an argument of Alzer and Berg [1] to derive an integral representation of $b_n$ and then we get some new properties about $b_n$ at once.

## 2. Lemmas

In order to prove our main results we need the following two lemmas. A proof of Lemma 1 is given in [1] .

**Lemma 1.** Let

$$f(x) = (x+1)[e - (\frac{1+x}{x})^x] \qquad (x > 0). \tag{2.1}$$

Then we have

$$f(x) = \frac{e}{2} + \frac{1}{\pi}\int_0^1 \frac{s^s(1-s)^{1-s}sin(\pi s)}{x+s}ds. \tag{2.2}$$

The next lemma can be found in [5].

**Lemma 2.** Let $h(s)$ is a continuous function on [0,1]. Then

$$\lim_{n\to\infty} n\int_0^1 s^n h(s)ds = h(1). \tag{2.3}$$

## 3. Results

The main result of this paper which incorporate the announced new properties of the sequence $\{b_n\}_{n\geq 1}$ in the following.

**Theorem 1.** Let $\{b_n\}_{n\geq 1}$be the sequence defined by (1.1) , and let

$$g(s) = \frac{1}{\pi}s^s(1-s)^{1-s}sin(\pi s).$$

Then

$$b_n = \frac{1}{e}\int_0^1 g(s)(1-s)^{n-2}ds = \frac{1}{e}\int_0^1 g(s)s^{n-2}ds, \tag{3.1}$$

$$0 < b_n \leq \frac{1}{n(n+1)} \qquad (n = 1, 2, \cdots), \tag{3.2}$$

$$b_{n+1} < b_n \qquad (n = 1, 2, \cdots), \tag{3.3}$$

$$\lim_{n\to\infty}\frac{b_{n+1}}{b_n} = 1. \tag{3.4}$$

*Proof.* Let

$$t = \frac{1}{1+x} \qquad (x > 0),$$

$$f(t) = e - (1-t)^{1-1/t}. \tag{3.5}$$

From (1.1) we obtain

$$b_n = \frac{f^{(n)}(0)}{n!e} \qquad (n = 1, 2, \cdots). \tag{3.6}$$

On the other hand , by Lemma 1 , we have

$$f(t) = \frac{e}{2}t + \int_0^1 g(s)\frac{t^2}{1+(s-1)t}ds, \tag{3.7}$$

where $0 \leq t \leq 1$ .
Differentiation gives

$$f'(0) = \frac{e}{2},$$

$$f^{(n)}(0) = n! \int_0^1 g(s)(1-s)^{n-2} ds \qquad (n = 2, 3, \cdots). \tag{3.8}$$

So that (3.6) and (3.8) yield

$$b_1 = \frac{1}{2},$$

$$b_n = \frac{1}{e} \int_0^1 g(s)(1-s)^{n-2} ds = \frac{1}{e} \int_0^1 g(s) s^{n-2} ds \qquad (n = 2, 3, \cdots), \tag{3.9}$$

where the last step uses the change of variable $s = 1 - t$. Which prove (3.1).
The relation (3.2) and (3.3) follow immediately from (3.9) and (1.1) .
Now , we prove (3.4). Since $g(1) = 0$ , we cannot directly use (3.9) to calculating this limit by Lemma 2.
Integrating by parts , we find from (3.9)

$$b_n = -\frac{1}{(n-1)e} \int_0^1 h(s) s^{n-1} ds \qquad (n = 2, 3, \cdots), \tag{3.10}$$

where

$$h(s) = s^s (1-s)^{1-s} [cos(\pi s) - \frac{sin(\pi s)}{\pi} \ln \frac{1-s}{s}].$$

Noting

$$\lim_{s \to 0^+} \frac{sin(\pi s)}{\pi} \ln \frac{1-s}{s} = 0,$$

$$\lim_{s \to 1^-} \frac{sin(\pi s)}{\pi} \ln \frac{1-s}{s} = 0,$$

from Lemma 2 and (3.9) , we have

$$\begin{aligned} &\lim_{n \to \infty} \frac{b_{n+1}}{b_n} \\ &= \lim_{n \to \infty} \frac{(n-1) \int_0^1 h(s) s^n ds}{n \int_0^1 h(s) s^{n-1} ds} \\ &= \lim_{n \to \infty} \frac{(n-1)^2}{n^2} \frac{(n) \int_0^1 h(s) s^n ds}{(n-1) \int_0^1 h(s) s^{n-1} ds} \\ &= \frac{h(1)}{h(1)} = \frac{-1}{-1} = 1. \end{aligned}$$

This completes the proof of Theorem 1. □

Remark That relation (3.1) can be rewritten as follows:

$$\int_0^1 g(s)(s)^{n-2} ds = \int_0^1 g(s)(1-s)^{n-2} ds = e b_n \qquad (n = 2, 3, \cdots).$$

which means that they are extension of [1,Lemma2]. For example

$$\int_0^1 g(s) ds = e b_2 = \frac{e}{24}, \qquad \int_0^1 g(s) s ds = e b_3 = \frac{e}{48},$$

and

$$\int_0^1 \frac{1}{s} g(s)ds = \int_0^1 \frac{1}{1-s} g(s)ds$$
$$= \int_0^1 (1 + s + s^2 + \cdots) g(s)ds$$
$$= e \sum_{n=2}^{\infty} b_n = e \sum_{n=1}^{\infty} b_n - eb_1$$
$$= e(1 - \frac{1}{e}) - \frac{e}{2} = \frac{e}{2} - 1.$$

## References


[1] H. Alzer, C. Berg , *Some classes of completely monotonic functions*, Ann. Acad. Sci. Fennicae, 27 (2002),445-460.

[2] M. Gyllenberg, P. Yan, On a conjecture by Yang, J. Math. Anal. Appl. 264 (2001) 687–690.

[3] G.H. Hardy, J.E. Littlewood and G. Polya, Inequalities, Cambridge Univ. Press, London, 1952.

[4] C. Mortici , Refinements of some bounds related to the constant $e$, Miskolc Math. Notes, 12 (2011), no. 1, 105-111.

[5] W. Trench, *Introduction to Real Analysis*, Prentice Hall ,2002.

[6] X.Yang, On Carleman's inequality, J. Math. Anal. Appl., 253 (2001), 691–694.

[7] X. Yang , *Approximations for constant e and Their Applications*, J.Math.Anal.APPL.262, 651-659 (2001).

[8] H. Yue, *A strengthened Carleman's inequality*, Comm. Math. Anal., 1 (2006) ,no. 2, 115-119.



College of Mathematics and Informatics, Henan Polytechnic University,, Jaozuo City, Henan 454000, P.R. China.

*E-mail address*: huu3y2@163.com

Valahia University of Târgovişte, Department of Mathematics, Bd. Unirii 18, 130082 Târgovişte, Romania

*E-mail address*: cmortici@valahia.ro